

ESTHETIC CROWN LENGTHENING WITH ALTERED PASSIVE ERUPTION

PATIENT NAME: Mr. H. M.

AGE: 21 years

GENDER: M

OPD NO: 212467

DIAGNOSIS: Generalized Dental Biofilm-induced gingivitis in an intact periodontium with tooth related factors with altered passive eruption type 1B or APE-I-AAE with 11, 12, 13, 21, 22, and 23

TREATMENT DONE: Crown lengthening procedure with maxillary anterior teeth

ESTHETIC CROWN LENGTHENING WITH ALTERED PASSIVE ERUPTION



FRONTAL VIEW



RIGHT LATERAL VIEW



LEFT LATERAL VIEW



IOPA WITH 21 AND 22



**INTERNAL BEVEL
INCISION WITH NO. 15
BLADE FOLLOWED BY
SULCULAR INCISION AND
VERTICAL RELEASING
INCISION GIVEN**



**FULL THICKNESS
MUCOPERIOSTEAL FLAP
REFLECTED**

ESTHETIC CROWN LENGTHENING WITH ALTERED PASSIVE ERUPTION



**ALVEOLAR BONE AT THE
LEVEL OF CEJ**



**AFTER OSTEOETOMY CEJ
IS VISIBLE**



**SUTURING THE FLAP 2-4
MM APICAL TO THE BONE
CREST**



COE PACK PLACED



**7 DAYS AFTER SUTURE
REMOVAL**



**15 DAYS AFTER SUTURE
REMOVAL**

ESTHETIC CROWN LENGTHENING WITH ALTERED PASSIVE ERUPTION



1 MONTH POST-OPERATIVE



**3 MONTH POST-OPERATIVE
AND WITH PROSTHESIS
WRT #21**