

## DEPTMENT OF ORAL MEDICINE AND RADIOLOGY

38 yr male patient visited to the department with complaint of burning sensation while having spicy food. Patient had habit history of tobacco chewing since 15 yrs on clinical evaluation it was reported that a homogenous white keratotic patch present on right buccal mucosa. Which is non scrapable and does not disappears on streaching. Provisional diagnosis was made as leukoplakia. patch type lichen planus given in differential diagnosis. Patient was advise to quit habit and ask them to apply kenacort ointment for 7 days and also avised antioxidant tablet s.m. fibro for 1 month. Pt had recalled after 1 month.





A 78 year old male pt reported to the department with chief complaint of burning sensation in left and right buccal mucosa. Patient gave history of tobacco chewing habit since, 8 year. On clinical examination it was noted that a grayish white keratotic appearance is seen with melanin pigmentation in background. The lesion is non scrapable and non reducible. Provisional diagnosis was made as reticular lichen planus. patient advices to apply kenacort ointment over the lesion 2-3 times per day



A 28 year female pt reported to the hospital with chief complaint of ulcer on lip. Patient gave history of burning sensation with aggravate while eating spicy food.

On examination a pseudo membranous slough was noted on labial mucosa surrounded by erythematous border. ulcer is having sloping borders and non indurated base.

Patient was advised to avoid hot and spicy food and instructed to apply lignocaine gel 2-3 times per day for 7 days. and multivitamin tablet such zincovit for 15 days.